

Many-body Systems Interacting via a Two-body Random Ensemble: average
energy of each angular momentum

Y. M. Zhao^{a,c}, A. Arima^b, and N. Yoshinaga^a

^a Department of Physics, Saitama University, Saitama-shi, Saitama 338 Japan

^b The House of Councilors, 2-1-1 Nagatacho, Chiyodaku, Tokyo 100-8962, Japan

^c Department of Physics, Southeast University, Nanjing 210018 China

In this paper, we discuss the regularities of energy of each angular momentum I averaged over all the states for a fixed angular momentum (denoted as $\bar{E}_I$'s) in many-body systems interacting via a two-body random ensemble. It is found that $\bar{E}_I$'s with $I \sim I_{min}$ (minimum of I) or I_{max} have large probabilities (denoted as $\mathcal{P}(I)$) to be the lowest, and that $\mathcal{P}(I)$ is close to zero elsewhere. A simple argument based on the randomness of the two-particle cfp's is given. A compact trajectory of the energy $\bar{E}_I$ vs. $I(I+1)$ is found to be robust. Regular fluctuations of the $\mathcal{P}(I)$ (the probability of finding I to be the ground state) and $\mathcal{P}(I)$ of even fermions in a single- j shell and boson systems are found to be reverse, and argued by the dimension fluctuation of the model space. Other regularities, such as why there are 2 or 3 sizable $\mathcal{P}(I)$'s with $I \sim I_{min}$ and $\mathcal{P}(I) \ll \mathcal{P}(I_{max})$'s with $I \sim I_{max}$, why the coefficients C defined by $\langle \bar{E}_I \rangle = CI(I+1)$ is sensitive to the orbits and not sensitive to particle number, are found and studied for the first time.

PACS number: 05.30.Fk, 05.45.-a, 21.60Cs, 24.60.Lz

1 Introduction

The discovery that the 0^+ dominance in ground states (0 g.s.) [1] of even fermion systems can be obtained by using a two-body random ensemble (TBRE) brought the physics of many-body systems interacting via a TBRE, such as the origin of this 0^+ dominance and its physical implications into a sharp focus [2-18]. However, a sound understanding of this 0^+ dominance is not yet available, though quite a few approaches were suggested. In this paper we discuss many-body systems interacting via a TBRE hamiltonian in another context: regularities of averaged energy (denoted as $\bar{E}_I = \frac{1}{dim_I} \sum_{\beta} E_{I\beta}$, where β represents additional quantum numbers of angular momenta I states and dim_I is the dimension of I states) for each angular momentum I .

Our study of $\bar{E}_I$ originates from the difficulties [16, 17, 18] that we met in studying the $P(I)$, the probability of finding angular momentum I to be the ground state. In Ref. [18] we studied the $P(I)$'s in two steps. The first step is to study the $\mathcal{P}(I)$'s, the probabilities of $\bar{E}_I$'s being the lowest. It is much simpler to understand the behavior of $\mathcal{P}(I)$'s than to understand the behavior of $P(I)$'s, because the $\bar{E}_I$'s are always linear combinations of two-body matrix elements, and the approach of [17] is applicable to predict the $\mathcal{P}(I)$'s of a randomly interacting many-body system. The second step is to study the features of width for each I . Unfortunately, such a picture is not applicable to odd number of fermions. The reason is that the correlation between states is also essential to explain the distribution of $P(I)$'s. Here correlation is an antonym of independence, it refers to, e.g., for fermions in a single- j shell, the state with $I_{max} - 2$ is very likely the first excited state when the $I = I_{max}$ state is the g.s.. It is not enough to consider only the statistical behavior of energy levels, especially for systems with few parameters (e.g., sd boson systems, fermions in a single- j shell).

Although we did not successfully relate the $\mathcal{P}(I)$'s with $P(I)$'s in [18], regularities of the $\mathcal{P}(I)$'s studied in [18] are very interesting and worthy of further study, as will be shown in this paper.

This paper is organized as follows: In Sec. 2, we present a few typical examples of $\mathcal{P}(I)$'s, which show that the $\bar{E}_I$'s with $I \sim I_{min}$ (minimum of I) or $I \sim I_{max}$ have large probabilities (denoted as $\mathcal{P}(I)$) to be the lowest, and that $\mathcal{P}(I)$ is close to zero elsewhere. A plausible understanding is given in terms of distributions of two-body coefficients of fractional parentage. In Sec. 3, we discuss a robust correlation between $\bar{E}_I$'s and $I(I+1)$ for all cases that we checked: fermions in a single- j or a two- j shell, d -, sd -, and sdg -boson systems. Furthermore, we define a “moment of inertia”, $\mathcal{J}$, by $\mathcal{J} = \frac{1}{2}I(I+1)/\langle\bar{E}_I\rangle$, and find that $\sqrt{\mathcal{J}}$ is proportional to j of fermions in a single- j shell (and d - and sd -boson systems), and almost independent of particle number n . Similar behavior is noticed in sdg -boson systems and fermion systems in two- j shell. Here, we find a regular even-odd staggering of both $\mathcal{P}(I)$'s and $P(I)$'s of even fermions in a single- j shell, and d -, sd -, and sdg -boson systems, which is argued by a regular odd-even staggering of dimension of the model space. One apparent observation that there are 2 or 3 sizable $\mathcal{P}(I)$'s with $I \sim I_{min}$ and only one large $\mathcal{P}(I_{max})$ with $\mathcal{P}(I) \ll \mathcal{P}(I_{max})$ ($I \sim I_{max}$) is interpreted as a reflection of fluctuations of $\bar{E}_I$ in each run of a TBRE hamiltonian. In Sec. 4 we reproduce the relation of $\bar{E}_I$'s and $I(I+1)$ and explain the features of $\mathcal{J}$'s for fermions in a single- j shell by assuming randomness of two-body cfps. A summary of this work will be given in Sec. 5.

2 Typical examples of $\mathcal{P}(I)$'s

In Fig. 1 we plot the $\mathcal{P}(I)$'s for a few very different systems: 4 fermions in a single- j shell ($j = \frac{15}{2}$), 6 fermions in a two- j ($2j_1, 2j_2$)=(11,7) shell, a 6- sdg -boson system, and 5 fermions in a single- j shell ($j = \frac{9}{2}$). The two-body interaction parameters are taken as the two-body random ensemble (TBRE), as most of previous works [2-18]. All the $\mathcal{P}(I)$'s are obtained by 1000 runs. We show in Fig. 1 typical examples among many cases that we have checked: 4, 5, and 6 valence fermions in a single- j shell up to $j = \frac{31}{2}$, both even or odd number of fermions ($n = 4$ to 9) in two- j shells with $(2j_1, 2j_2)$ =(7,5), (11,3), (11, 5), (11,7), (11,9), (13,9), d -boson systems with n_d changing from 3 to 45, sd -boson systems with n changing from 4 to 17, and sdg -boson systems with $n = 4, 5, 6$. It is noted that the $\mathcal{P}(I)$'s are large if $I \sim I_{min}$ or I_{max} ; they are close to zero elsewhere. This feature is general for all cases. It is noted that a displacement on the TBRE produce only a constant shift on the ground state energy. Thus it is robust regardless of the displacement, while $P(I)$ is quite sensitive to the displacement [7].

Now we give an understanding on this observation. Let $\bar{E}_I = \sum_{k_0} \bar{\alpha}_I^{k_0} G_{k_0}$, where G_{k_0} 's are two-body matrix elements determined randomly (TBRE parameters), and $\bar{\alpha}_I^{k_0}$'s are coefficients calculated in terms of two-particle cfp's. For the sake of simplicity, here we restrict ourselves to fermions in a single- j shell but a generalization to other cases, we believe, is straightforward. In a single- j shell, k_0 is uniquely labeled by J [17]. We use G_J and $\bar{\alpha}_I^J$ instead of G_{k_0} and $\bar{\alpha}_I^{k_0}$ for a single- j shell. As a specific example, we discuss the case of $j = \frac{9}{2}$ with 4 fermions below.

Using the coefficients $\bar{\alpha}_I^J$, we predict the $\mathcal{P}(I)$'s by using integrals similar to Eq. (7) of Ref. [17], without diagonalizing a TBRE hamiltonian. The predicted $\mathcal{P}(I)$'s for 4 fermions in a $j = \frac{9}{2}$ shell are listed in the column "pred1." in Table I. It is easy to notice that the $\mathcal{P}(I)$ is large if $\bar{E}_I$ has one or more largest (and the smallest) $\bar{\alpha}_I^J$'s for different I 's but a fixed G_J . The predicted $\mathcal{P}(I)$'s by the simple approach proposed in [16], $\mathcal{P}(I) = \mathcal{N}_I / N_m$, are listed in the column "pred2." of Table I, where $N_m = 2N - 1$, and $\mathcal{N}_I$ is the number of angular momenta I being either the smallest

or the largest eigenvalues with only one of two-body matrix elements switched on. The $\mathcal{P}(I)$'s obtained by diagonalizing a TBRE hamiltonian (1000 runs) are given in the column "TBRE". It is seen that all predicted $\mathcal{P}(I)$'s are reasonably consistent with those obtained by diagonalizing a TBRE hamiltonian.

It is interesting to ask why the largest $\bar{\alpha}_I^J$'s (for different I and fixed J) appear in $\bar{E}_I$'s with $I \sim I_{min}$ or I_{max} . Our interpretation is as follows.

First, $\bar{\alpha}_I^J$'s are the largest if $I \sim I_{min}$ (small J cases) or I_{max} (large J cases) by assuming that the two-particle cfp's, and hence the $\alpha_{I\beta\beta'}^J$'s are **randomly** given. This assumption was found [18] to work well except at the edge of angular momenta I . As a specific example, we show the distribution of cfp's and $\alpha_{I\beta\beta'}^J$'s for 4 fermions in a $j = \frac{31}{2}$ shell in Fig. 2. It is interesting to note that both the cfp's and $\alpha_{I\beta\beta'}^J$'s seem to follow a Poissonian distribution, the origin of which is not known.

We thus define d_I^J/D_I , a ratio between the dimension of two-particle cfp's corresponding to interaction G_J of angular momentum I state and the total dimension of the two-particle cfp's of angular momentum I state. The $d_I^J/D_I \times \frac{1}{2}n(n-1)$ is then a reasonable estimate of $\bar{\alpha}_I^J$ [17, 19] if one assumes the two-particle cfp's are "uniformly" distributed in the D_I components. It is easy to realize that $\bar{\alpha}_I^J$'s for states with small J and $I \sim I_{min}$ are larger than those of other states because d_I^J/D_I of small J and $I \sim I_{min}$ states are mostly larger. Similar statements are applicable to the $\bar{\alpha}_I^J$'s of the states with $I \sim I_{max}$ and large J . To exemplify this argument, in Table II we list d_I^J and D_I of 5 fermions in a $j = \frac{15}{2}$ shell, where the d_I^J/D_I with $J < j - \frac{1}{2}$ and $I \sim I_{min}$ are mostly larger than that of states elsewhere, and the d_I^J/D_I with $J > j - \frac{1}{2}$ and $I \sim I_{max}$ are larger than that of other states.

Second, the fluctuations of $\bar{\alpha}_I^J$ with $I \sim I_{min}$ or I_{max} are larger than those with medium I . In all cases, including boson systems, even or odd number of fermions in a single- j or a many- j shell, the dimension of two-particle cfp's and that of the model space with $I \sim I_{min}$ or I_{max} are smaller than those with medium I , which suggests that the fluctuations of the $\bar{\alpha}_I^J$ with $I \sim I_{min}$ or I_{max} are much larger than those elsewhere if it is assumed that the two-particle cfp's are "random".

Therefore, the statistical properties of two-particle cfp's and the dimension of two-particle cfp's, and the property of the shell model dimension accounts for the facts why there are always many $\bar{\alpha}_I^J$'s, which are the largest (or the smallest) for different I and fixed J , in $\bar{E}_I$'s with $I \sim I_{min}$ or I_{max} .

A “staggering” of $\mathcal{P}(I)$'s is noticed for boson systems and even fermion in a single- j shell: the $\mathcal{P}(I)$'s of odd I are larger than their even neighbors with very few exceptions. In contrast to this systematics, it was noticed [9, 17] that the $P(I)$'s for even I in these systems are larger than their odd neighbors. These two “reversed” regularities are explained as a reflection of a regular change in dimension of the model space below: For boson systems and even fermions in a single- j shell, the dimension of the model space for even I is systematically and relatively larger than their odd I neighbors. The $\mathcal{P}(I)$'s are determined by the coefficients $\bar{\alpha}_I^k$. The relatively larger dimensions of the model space for even I 's equivalently lead to relatively smaller fluctuations of $\bar{\alpha}_I^k$ of even I 's, and the $\bar{\alpha}_I^k$ are relatively “medium” compared with their odd- I neighbors. This gives relatively larger $\mathcal{P}(I)$'s for odd I 's. On the other side, the $P(I)$'s are determined by the largest eigenvalues with one of the two-body matrix elements switched on. The larger dimension of the model space for even I states equivalently produces a relatively larger maximum (denoted as E_I^0) of E_I 's for even I 's than their odd neighbors, where the E_I^0 's are obtained by diagonalizing two-body matrix elements N times, in each time only one of $G_k = 1$ and the other interactions are switched off.

3 A regularity of averaged energies of fixed angular momenta

If one examines the spin ordering of the average energies $\bar{E}_I$'s, it is easy to notice that if the angular momentum I of the lowest $\bar{E}_I$ belongs to the case of $I \sim I_{min}$ (I_{max}) in **one** run of the TBRE, very likely the average energy $\bar{E}_I$ increases (decreases) with the total angular momentum I , and is proportional to $I(I+1)$ on average. The probability of $\bar{E}_I$ being the lowest with “medium” I is close to zero.

Let $\langle \bar{E}_I \rangle_{min}$ ($\langle \bar{E}_I \rangle_{max}$) be a quantity obtained by averaging $\bar{E}_I$ over the cases with only $I \sim I_{min}$ ($I \sim I_{max}$) g.s. for the ensemble used. We find that both $\langle \bar{E}_I \rangle_{max}$ and $\langle \bar{E}_I \rangle_{min}$ are proportional to the $I(I+1)$, similar to a “rotational” spectra.

In Fig. 3 we show $\langle \bar{E}_I \rangle_{min}$ vs. $I(I+1)$ for a d -boson system with $n = 20$, a sd -boson system with $n = 10$, and a system with 4 fermions in a single- j ($j=17/2$) shell, and a system with 6 fermions in a two- j shell. For the sake of simplicity, below we introduce “moment of inertia” $\mathcal{J}$, defined by a optimal constant $(2C)^{-1}$, where C is the coefficient defined in $\langle \bar{E}_I \rangle_{max}$ ($\langle \bar{E}_I \rangle_{min}$) = $CI(I+1)$.

In the pioneering paper [1] by Johnson et al, they showed in their Fig. 4 that the average excitation energy follows roughly a function proportional to $I(I+1)$, but the energy they referred to are average energy of yrast and even I states, therefore it is different from the $\langle \bar{E}_I \rangle_{min}$ in this paper. It is noted that we consider both even I and odd I in our “rotation”.

Prior to this work, based on geometric chaoticity, Mulhall, Zelevinsky, and Volya [9] elaborated a formula below:

$$E_I = \text{Const}_1 + \text{Const}_2 I(I+1) + o(I(I+1)). \quad (1)$$

Those authors discussed the $P(I)$'s by using the above formula for fermions in a single- j shell. It seems, however, that the E_I in their formula should correspond to our $\langle E_I \rangle$. The features predicted by [9], such that the minimum and the maximum of angular momenta I are favored to be the ground states with around 50% for each (sum of all $\mathcal{P}(I)$'s with $I \sim I_{min}$ or I_{max}), and that the $\langle \bar{E}_I \rangle_{min}$'s (and $\langle \bar{E}_I \rangle_{max}$)

behave like a rotor, seem more appropriate to describe the $\mathcal{P}(I)$'s, $\langle \bar{E}_I \rangle_{min}$ and $\langle \bar{E}_I \rangle_{max}$, rather than the $P(I)$'s and the energy obtained by averaging yrast energy over the cases of 0 g.s. or I_{max} g.s. in the ensemble. The “equilibrium” energies obtained in the statistical way of [9] may correspond to the energies averaged over all states with given angular momenta I 's, not the yrast ones. In fact, the $I(I+1)$ behavior in average yrast energy defined in ref. [9] was not well confirmed.

In another recent work [7], the spectroscopy of $\bar{E}_I$ was checked but this pattern was not noticed.

It is noted that one should not confuse $\bar{E}_I$'s with Bethe expression of level densities [20] which is based on a Fermion gas approach. First, the $\bar{E}_I$'s are always Gaussian and $\bar{E}_I$'s obtained by averaging over a TBRE should be zero. The $I(I+1)$ behavior appears when one divides a TBRE into two cases— $I \sim I_{min}$ g.s. and $I \sim I_{min}$ g.s.—and calculates $\langle \bar{E}_I \rangle_{min}$ and $\langle \bar{E}_I \rangle_{max}$. Second, it is emphasized that an $I(I+1)$ behavior of $\bar{E}_I$ in the Bethe expression of level density and that in $\langle \bar{E}_I \rangle_{min}$ (or $\langle \bar{E}_I \rangle_{max}$) are completely different. For example, $\mathcal{J}$ in Bethe expression changes with particle number but $\mathcal{J}$ in this work, as will be shown, is not sensitive to particle number n . Furthermore, systems discussed in this paper can be very simple, and those described by Bethe expression require complexity in energy levels so that one needs statistical approach for level densities. The $I(I+1)$ behavior discussed in this paper was not reported in previous works.

In Fig. 4 we plot C for fermions in a few single- j or two- j shells. The results indicate:

1. C is almost independent of particle number n for all systems.
2. C of fermions in a single- j shell is quite sensitive to orbits (labeled by j), that of fermions in a two- j shell is sensitive to the larger j value. The coefficient C decreases with the larger j if the smaller j 's are the same. For instance, coefficient C of $(2j_1, 2j_2) = (7, 5)$, that of $(2j_1, 2j_2) = (9, 5)$ and that of $(2j_1, 2j_2) = (11, 5)$ decrease successively.
3. For fermions in a two- j shell, the difference between coefficients C of systems

with the same j (larger) and $j_2 \ll j_1$ is very small. For example, the coefficients C for shells $(2j_1, 2j_2) = (11, 1), (11, 3)$ are very close.

Without presenting details we mention that the C determined by $\langle \bar{E}_I \rangle_{min} = CI(I+1)$ is close to that by $\langle \bar{E}_I \rangle_{max} = CI(I+1)$ for all examples that we have checked. This could be easily understood by the symmetry of the ensemble that we use.

An empirical relation between $\mathcal{J}$ and j is summarized in Fig. 5. For fermions in a single- j shell, d boson systems and sd boson systems, $\mathcal{J}$ is fitted by a trajectory of $\sqrt{\mathcal{J}} = 1.42j$ (we take $j = 2$ for d boson and sd boson systems); for sdg bosons and fermions in a two- j shell, there seems a slight shift from the trajectory $\sqrt{\mathcal{J}} = 1.42j$, where we take $j^2 = j_1^2 + j_2^2$. It is noted that one cannot, though the Eq. (7) of [9] seems to indicate a correct behavior of $\langle \bar{E}_I \rangle$, obtain this simple systematics of $\mathcal{J}$ based on their formulas. The $\mathcal{J}$ of [9] is very different from that given here. Therefore, this regularity cannot be explained by previous schemes such as [9] and deserves further studies.

Another feature of $\mathcal{P}(I)$'s is that the $\mathcal{P}(I_{max})$'s are always quite "stable" ($\sim 28\text{-}35\%$), while the $\mathcal{P}(I_{max} - 2)$'s of fermions in a single- j shell and boson systems, and $\mathcal{P}(I_{max} - 1)$'s of fermions in a many- j shell, is drastically smaller than those of the I_{max} state(s), though still sizable ($\sim 5 - 15\%$), and that there may be 2 or 3 sizable $\mathcal{P}(I)$'s for the cases of $I \sim I_{min}$, and the $\mathcal{P}(I_{min})$'s are not *always* larger than other $\mathcal{P}(I)$'s (with $I \sim I_{min}$).

Now we are able to explain the asymmetry of $\mathcal{P}(I)$ by using the fluctuations of $\bar{E}_I$. The $\bar{E}_I$'s are proportional to $I(I+1)$ but with fluctuations in each run of a TBRE hamiltonian. Because $(\bar{E}_{I+1} - \bar{E}_I)$ is small if I is small and large if I is large, the probabilities to change order of $\bar{E}_I$'s for $I \sim I_{min}$ due to the fluctuations of $\bar{E}_I$ is much larger than those for $I \sim I_{max}$. That is why there are 2 or 3 sizable and comparable $\mathcal{P}(I)$'s with $I \sim I_{min}$ but only one large $\mathcal{P}(I)$'s with $I \sim I_{max}$ and the $\mathcal{P}(I_{max} - 1) \ll \mathcal{P}(I_{max})$.

4 A scenario of $\langle \bar{E}_I \rangle \sim I(I+1)$ relation

It is more convincing if one “reproduces” the $\langle \bar{E}_I \rangle I(I+1)$ relation by the above assumption of randomness of two-body cfp’s. Below we first show by one specific case that $\langle \bar{E}_I \rangle$ vs. $I(I+1)$ for fermions in a single- j shell, then we present a simple approach to evaluate $\mathcal{J}$ of fermions in a single- j shell. Note that a simple approach to evaluate the $\mathcal{J}$ of fermions in a many- j shell is not yet available.

As indicated in Fig. 4, $\mathcal{J}$ is not sensitive to particle number n . Thus we study only 4-fermion systems. We concentrate on states with $I \geq 2j - 1$, where the dimension of two-particle cfp’s is very simple:

$$D_I = \mathcal{L}(\mathcal{L} + 1)/2, \\ d_I^J = \begin{cases} \mathcal{L} + J/2 + (1 - 2j)/2, & \text{if } 2j \leq \mathcal{L} + J + 1 - 2j \\ 0 & \text{otherwise} \end{cases} ; \quad (2)$$

where $\mathcal{L} = 2j - [I/2]$. Suppose that the two-particle cfp’s are uniformly distributed, and that the number of two-particle cfp’s are large enough for a statistical prediction. We have $\bar{\alpha}_I^J = d_I^J / D_I \times n(n-1)/2$. It was checked in [18] that the $\bar{\alpha}_I^J$ ’s predicted in this way is well consistent with those by diagonalizing matrices if the dimension of two-particle cfp’s is quite large.

Next we use these predicted $\bar{\alpha}_I^J$ ’s and random two-body interaction parameters (TBRE), G_J ’s, to evaluate the averaged energies of given angular momenta. As an example we use $j = 15/2$ and $n = 4$. The results are shown in Fig. 6a). It is seen that the averaged energies, $\langle \bar{E}_I \rangle$ ’s, are approximately proportional to $I(I+1)$.

Now let us introduce an approximation of $\langle G_J \rangle_{min}$ ’s by only 3 requirements for 4 fermions in a single- j shell. The first requirement is that $\sum_j \langle G_J \rangle_{min} \sim 0$, and the second requirement is that $\langle G_{J_{max}} \rangle_{min} = 0.7$, $\langle G_{J_{max}-k} \rangle_{min} \sim 0.3$ with $k = 2, 4$. The last requirement is $\langle G_J \rangle_{min} \sim -0.3$ when $J < 8$. Other $\langle G_J \rangle_{min}$ is quite small in magnitude. Those requirements are based on our calculations of fermions in a single- j shell. By using these $\langle G_J \rangle$ ’s and $\bar{\alpha}_I^J$ ’s, we obtain Fig. 6b). It is seen that the results are very similar: The $\mathcal{J}$ obtained in Fig. 6a) is 9.96, that obtained in

Fig. 6b) is 11.18, and that in Fig. 5, which is obtained by diagonalizing a TBRE hamiltonian, is 10.30. Therefore, we show clearly by this example that the $I(I+1)$ behavior of $\bar{E}_I$'s, at least for 4 fermions in a single- j shell, is a reflection of a random distribution of the two-particle cfp's.

Below we estimate $\mathcal{J}$, and thereby giving a simple argument on the behavior of $\mathcal{J}$ of fermions in a single- j shell. We take that $\langle \bar{E}_{I_{min}} \rangle \sim 0$, and $I_{min}(I_{min}+1) \sim 0$. We assume that $\langle \bar{E}_{I_{max}} \rangle \sim \alpha_{I_{max}}^{J_{max}} \langle G_{J_{max}} \rangle$. For the sake of simplicity, we take that $\langle G_{J_{max}} \rangle = 0.7$ for $n = 4$ in all single- j shells. Then we have that for $n = 4$:

$$\alpha_{I_{max}}^{J_{max}} \langle G_{J_{max}} \rangle \sim \frac{1}{2\mathcal{J}}(4j-6)(4j-5) \sim \frac{8}{\mathcal{J}}j^2. \quad (3)$$

From [16], it is seen that the $\alpha_{I_{max}}^{J_{max}}$ of $n = 4$ saturates quickly at $\frac{29}{8}$. Then the left hand side $\sim \frac{29}{8} \times 0.7 \sim 2.54$. We find that $\sqrt{\mathcal{J}} \sim 1.77j$, a bit larger than (but close to) the $\sqrt{\mathcal{J}}$ obtained in Fig. 5 ($\sqrt{\mathcal{J}} = 1.42j$). The $\mathcal{J}$ in Eq. (3) is over-estimated because we neglect the contributions from $G_{J_{max}-2}$ and $G_{J_{max}-4}$ in this simple phenomenology.

The $\mathcal{J}$ of d - and sd -boson systems may be estimated in a similar way. For example, $c_4 - c_2 > 0$ gives $\bar{E}_I$ with $I \sim 0$ ground states. The $c_4 - c_2 > 0$ has an average around 0.88. From Eqs. (10-11), one has that $\mathcal{J} \sim 7/(c_4 - c_2) \sim 7.95$, or $\sqrt{\mathcal{J}} \sim 2.82$, consistent with that in Fig. 5 (2.67).

5 Summary and discussion

To summarize, we have presented in this paper for the first time two main robust regularities of many-body systems interacting via a two-body random ensemble: 1. The $\mathcal{P}(I)$'s, probabilities of average energies $\bar{E}_I$ of the angular momentum I states being the lowest for many-body systems, are large only and only if $I \sim I_{min}$ or I_{max} . 2. The $I(I+1)$ behavior of $\langle \bar{E}_I \rangle_{min}$ and $\langle \bar{E}_I \rangle_{max}$. Without detailed discussions it is noted in this paper that the regularities of $\mathcal{P}(I)$'s remain essentially the same if one takes other ensembles, such as the displaced random numbers, or random numbers which are only positive (or negative), while those of $P(I)$'s by using other ensembles may be completely different from those obtained by a TBRE hamiltonian. This suggests that the regularities of the $\mathcal{P}(I)$'s and $\langle \bar{E}_I \rangle$ are very **robust**.

We first propose an approach to interpret the regularity of $\mathcal{P}(I)$'s in terms of randomness of coefficients $\bar{\alpha}_I^k$, which are obtained from the two-particle cfp's and the approach developed in Ref. [17]. It is suggested that the dimension of two-particle cfp's and the that of the model space accounts for the large $\bar{\alpha}_I^k$ with $I \sim I_{max}$ and I_{min} .

The “staggering” patterns of $\mathcal{P}(I)$'s and $P(I)$'s are found to be reverse in boson systems and even fermions in a single- j shell: $P(I)$'s ($\mathcal{P}(I)$'s) with even I is systematically larger (smaller) than those of their odd I neighbors. These staggering patterns are interpreted in terms of a regular staggering in dimension of the model space. The asymmetry of distribution of $\mathcal{P}(I)$'s is explained by fluctuations of $\bar{E}_I$.

We provide a scenario of “reproducing” a compact trajectory of $\langle \bar{E}_I \rangle$ plotted against $I(I+1)$ by assuming the randomness of two-particle cfp's in the case of fermions in a single- j shell. We also propose a simple method to estimate $\mathcal{J}$ and a simple method to simulate the linear relation between $\sqrt{\mathcal{J}}$ of fermions in a single- j shell and the angular momentum of the orbit, j .

We therefore believe that the randomness of two-particle cfp's is the origin of all the observed regularities related to $\bar{E}_I$'s and $\mathcal{P}(I)$'s.

Finally, it is pointed out that the $\mathcal{P}(I)$'s and $P(I)$ discussed in our previous

papers [17, 18] are different quantities. For even systems the behavior of these two are accidentally similar. For odd- A systems, however, the $\mathcal{P}(I)$'s are very different from $P(I)$'s, which explicitly demonstrates that the I g.s. probabilities (and 0 g.s. dominance) cannot be explained by geometric chaoticity [9]. The 0 g.s. dominance is actually related to two-body matrix elements which have specific features [16].

Acknowledgement: One of us (YMZ) is grateful to Drs. S. Pittel, Y. Gono, Y. R. Shimizu, and R. Bijker for discussions and/or communications. This work is supported in part by the Japan Society for the Promotion of Science under contract No. P01021.

References

- [1] C. W. Johnson, G. F. Bertsch, D. J. Dean, Phys. Rev. Lett. **80**, 2749(1998).
- [2] C. W. Johnson, G. F. Bertsch, D. J. Dean, and I. Talmi, Phys. Rev. **C61**, 014311(1999).
- [3] C. W. Johnson, Rev. Mex. Fis. **45** **S2**, 25(1999).
- [4] R. Bijker, A. Frank, Phys. Rev. Lett. **84**, 420(2000); Phys. Rev. **C62**, 14303(2000).
- [5] D. Kusnezov, N. V. Zamfir, and R. F. Casten, Phys. Rev. Lett. **85**, 1396(2000).
- [6] M. Horoi, B. A. Brown, V. Zelevinsky, Phys. Rev. Lett. **87**, 062501(2001).
- [7] V. Velazquez, and A. P. Zuker, Phys. Rev. Lett. **88**, 027502(2002); Y. M. Zhao, and A. Arima, to be published.
- [8] R. Bijker, A. Frank, and S. Pittel, Phys. Rev. **C60**, 021302(1999).
- [9] D. Mulhall, A. Volya, and V. Zelevinsky, Phys. Rev. Lett. **85**, 4016(2000); Nucl. Phys. **A682**, 229c(2001); D. Mulhall, V. Zelevinsky, and A. Volya, nucl-th/0103069; V. Zelevinsky, D. Mulhall, and A. Volya, Yad. Fiz. **64**, 579(2001).
- [10] D. Kusnezov, Phys. Rev. Lett. **85**, 3773(2000); *ibid.* **87**, 029202 (2001); R. Bijker, and A. Frank, Phys. Rev. Lett. **87**, 029201(2001).
- [11] R. Bijker, and A. Frank, Phys. Rev. **C64**, (R)061303(2001).
- [12] R. Bijker, and A. Frank, Phys. Rev. **C65**, 044316(2002).
- [13] Lev Kaplan, Thomas Papenbrock, and Calvin W. Johnson, Phys. Rev. **C63**, 014307(2001).
- [14] S. Drozd and M. Wojcik, Physica **A301**, 291(2001).

- [15] J. Y. Zhang, N. V. Zamfir, R. F. Casten, and M. A. Caprio, Phys. Rev. **C64**, 017302(2001).
- [16] Y. M. Zhao, A. Arima, and N. Yoshinaga, nucl-th/112075; to be published.
- [17] Y.M. Zhao, and A. Arima, Phys. Rev. **C64**, (R)041301(2001).
- [18] A. Arima, N. Yoshinaga, and Y.M. Zhao, Eur. J. Phys. A **13**, 105(2002); N. Yoshinaga, A. Arima, and Y.M. Zhao, J. Phys. **G**, in press.
- [19] R. D. Lawson, *Theory of the Nuclear Shell Model* (Clarendon, Oxford, 1980).
- [20] A. Bohr, and B. Mottelson, *Nuclear structure*, Vol. I (Benjamin, Reading, 1964).

TABLE I. The coefficients $\bar{\alpha}_I^J$ for 4 fermions in a $j = \frac{9}{2}$ shell. Bold font is used for the largest $\bar{\alpha}_I^J$ are the largest and italic for the smallest $\bar{\alpha}_I^J$ for a given J . The probabilities in the column “pred1.” are obtained by integrals similar to Eq. (7) in Ref. [17], and those in the column “pred2.” are obtained by the approach discussed in Ref. [16]. The $\mathcal{P}(I)$ ’s in the last column “TBRE” (in %) are obtained by diagonalizing a TBRE hamiltonian 1000 runs. We take both the smallest and the largest $\bar{\alpha}_I^J$ when counting $\mathcal{N}_I$.

I	G_0	G_2	G_4	G_6	G_8	Pred1.(%)	pred2.(%)	$\mathcal{P}(I)$
0	0.80	0.35	1.74	2.11	1.01	11.97	11.1	10.2
2	0.30	1.39	1.45	<i>1.29</i>	1.56	14.51	22.2	15.4
3	0.00	0.36	2.28	2.63	<i>0.71</i>	28.17	33.3	28.9
4	0.20	1.07	1.38	1.91	1.44	1.74	0	1.7
5	0.00	1.00	1.59	1.84	1.57	0.30	0	0.6
6	0.20	0.79	1.50	1.58	1.93	0.22	0	0.3
7	0.00	1.20	1.09	1.40	2.31	3.44	0	3.2
8	0.30	0.48	1.05	1.82	2.36	0.03	0	0
9	0.00	0.17	1.33	2.12	2.38	0.01	0	0
10	0.00	0.70	0.69	1.41	3.21	6.76	0	8.7
12	0.00	<i>0.00</i>	<i>0.52</i>	1.69	3.78	32.64	33.3	31.0

TABLE II. The dimension of two-particle coefficients of fractional parentage (cfp's) in the case of single- $j = \frac{15}{2}$ with 5 fermions. The total dimension of two-body cfp's for different I and that of the shell model space are given in the two columns “cfp's” and “SM”, respectively. Note that the dimension distribution of two-body cfp's for fermions in a larger single- j shell or a many- j shell is similar. The $\mathcal{P}(I)$'s in the last column (in %) are obtained by diagonalizing a TBRE hamiltonian 1000 runs. Bold font is used for the d_I^J/D_I 's which give the largest, and italic font is used for those which are very comparable.

$2I$	G_0	G_2	G_4	G_6	G_8	G_{10}	G_{12}	G_{14}	cfp's	SM	$\mathcal{P}(I)$
1	0	2	<i>3</i>	4	5	<i>4</i>	3	3	24	2	18.4
3	1	3	<i>6</i>	9	9	<i>8</i>	7	5	48	4	12.8
5	1	<i>5</i>	<i>9</i>	12	13	<i>12</i>	10	8	70	6	12.6
7	1	7	<i>12</i>	15	17	<i>16</i>	13	11	92	8	3.6
9	2	8	<i>14</i>	18	19	<i>19</i>	17	13	110	9	1.5
11	2	10	<i>16</i>	20	22	<i>22</i>	20	16	128	11	1.4
13	2	11	<i>17</i>	21	24	<i>24</i>	22	19	140	11	0.1
15	3	11	17	23	25	<i>26</i>	25	22	152	13	0.1
17	2	11	18	23	26	<i>27</i>	26	23	156	12	0
19	2	11	18	23	27	<i>28</i>	27	25	161	13	0
21	2	9	17	23	26	<i>27</i>	28	27	159	12	0
23	2	9	16	22	26	<i>28</i>	28	27	158	12	0
25	1	8	14	20	25	<i>27</i>	28	28	151	11	0
27	2	7	13	19	23	<i>26</i>	28	28	146	11	0
29	1	6	11	16	21	<i>25</i>	27	28	135	9	0
31	1	6	10	14	19	<i>23</i>	26	28	127	9	0
33	1	4	8	12	16	<i>21</i>	25	27	114	7	0
35	1	4	7	10	14	<i>19</i>	23	26	104	7	0.1
37	0	3	5	8	12	<i>16</i>	21	25	90	5	0
39	1	2	4	7	10	<i>14</i>	19	23	80	5	0.4
41	0	2	3	5	8	<i>12</i>	16	21	66	3	0
43	0	2	2	4	7	<i>10</i>	14	19	57	3	0
45	0	0	1	3	5	<i>8</i>	12	16	45	2	2.0
47	0	0	1	2	4	<i>7</i>	10	14	38	2	2.3
49	0	0	0	1	3	<i>5</i>	8	12	29	1	5.5
51	0	0	0	1	2	<i>4</i>	7	10	24	1	10.3
55	0	0	0	0	1	2	4	7	14	1	28.9

Figure captions:

FIG. 1 Typical behaviors of the $\mathcal{P}(I)$'s: a) single- j ($j = \frac{15}{2}$) with 4 fermions, b) two- j shell $(2j_1, 2j_2) = (11, 7)$ with 6 fermions, c) 6-*sdg* bosons, d) single- j ($j = \frac{9}{2}$) with 5 fermions,

FIG. 2 Typical distribution of two-particle cfp's and $\alpha_{I\beta\beta'}^J$ of 4 fermions in a $j = 31/2$ shell and a medium angular momentum I ($=20$). It is seen that the distributions are close to Poissonian. The distributions of other I are similar unless $I \sim I_{min}$, or I_{max} , where the numbers of the two-particle cfp's and $\alpha_{I\beta\beta'}^J$ are small.

FIG. 3 Typical behavior of $\langle \bar{E}_I \rangle_{min}$ vs. $I(I+1)$. The $\langle \bar{E}_I \rangle_{min}$'s are obtained by averaging over all $\bar{E}_I$ given by diagonalizing a TBRE hamiltonian, with a requirement that $I \sim I_{min}$ g.s. a). 20-*d* bosons; b) 10-*sd* bosons; c) single- j shell with $j=17$ and $n = 4$, and d) a two- j shell with $j_1 = 5/2, j_2 = 7/2$, and $n = 4$. The reversed cases, i.e., the $I \sim I_{max}$ g.s., are very similar to the cases of $I \sim I_{min}$ g.s..

FIG. 4 Coefficients C (in $\langle \bar{E}_I \rangle_{min} = CI(I+1)$) of fermions in different shells. It is seen that C is not sensitive to particle number n , but rather sensitive to j values of the shell. Refer text for details.

FIG. 5 Correlation between $\sqrt{\mathcal{J}}$ and $j = \sqrt{\sum_i j_i^2}$. It is indicated that $\mathcal{J}$ is approximately proportional to $\sum_i j_i^2$ for *d*- and *sd*-bosons, and fermions in a single- j shell, with $\mathcal{J}$ of many- j shells and *sdg*-boson shifted very slightly to the right.

FIG. 6 A scenario of $\bar{E}_I \sim I(I+1)$. for 4 fermions in a single- j ($j = 15/2$) shell. The dot straight lines are used to guide the eyes. Only even I are plotted, because the predicted $\bar{E}_{I+1}$ (I is even) is the same as the predicted $\bar{E}_I$. Solid line is obtained by a linear fit. a) Solid squares are obtained by using predicted $\bar{\alpha}_I^J$ ($= d_I^J/D_I \times \frac{1}{2}n(n-1)$) and taking G_J 's to be TBRE. b) Solid squares are obtained

by using the same $\bar{\alpha}_I^J$ with a simple assumption of G_J . Refer to the text for details.




FIG. 1

YM ZHAO July/16th/01




FIG. 2




FIG. 3

FIG. 4





FIG. 5

Y.M.ZHAO, January 14th/2002






FIG. 6